

Stat310: Test 2

Answer the questions in the spaces provided on the question sheets. If you run out of room for an answer, continue on the back of the page. You have **80 minutes** and you may use one double-sided page of notes. You may not use your text book or a computer (calculators are fine). Please pledge on the last page.

Name: _____

1. Let P be a random variable that is 1 if someone smokes pot and 0 otherwise. $P \sim \text{Bernoulli}(p)$
 - (a) What does $P \sim \text{Bernoulli}(p)$ mean in this context? A person either smokes pot or doesn't, so where does the randomness come from? (1)

 - (b) We can estimate the true value of p by performing a survey and asking each participant whether or not they smoke pot. Let Q_i be 1 if person i smokes pot and 0 otherwise. Use the plug-in principle to write an estimate for p if $i = 1, \dots, n$. (2)

 - (c) Compute the variance of that estimator. (Hint: think about the distribution of the sum) (3)

Another way to estimate p is to use a randomised response question like we did in class. Instead of asking the participant directly, we can ask them to flip a coin and say yes if it's heads, and to answer truthfully otherwise. Let H be 1 for heads and 0 for tails, and let R be 1 if they answer yes to this question and 0 otherwise.

- (d) Fill in the blanks and then use that information to derive the joint distribution of H and R , and then the marginal distribution of R . (3)

$$P(H = 0) = \underline{\hspace{2cm}} \quad P(H = 1) = \underline{\hspace{2cm}}$$

$$P(R = 1|H = 0) = \underline{\hspace{2cm}} \quad P(R = 1|H = 1) = \underline{\hspace{2cm}}$$


	H = 0	H = 1
R = 0		
R = 1		

$$P(R = 0) = \underline{\hspace{2cm}} \quad P(R = 1) = \underline{\hspace{2cm}}$$

- (e) Give a transformation of P and H that produces R . (1)

- (f) If $\bar{R}_n = \sum_i^n R_i/n$, then we can estimate p with $2\bar{R}_n - 1$. Show that the expected value of this statistic is p . (2)

- (g) The following figure shows the variance of the two estimators for different values of true p . What do you notice? Is there any reason you would choose an estimator based on R instead of Q ? (3)



2. Let X_i be independent $\text{Poisson}(\lambda_i)$. (Therefore the mgf of X_i is $M_{x_i}(t) = e^{\lambda_i(e^t-1)}$).

- (a) Let $S_n = \sum_i^n X_i$. What is M_{S_n} , the mgf of S_n ? (3)

- (b) What is the distribution of S_n ? (2)

- (c) If $\lambda_i = \lambda$ and n is large, give a distribution (with parameters) that will closely approximate the distribution of S_n . (3)

(d) Suggest two estimators for λ using the plug-in principle. (2)

3. The joint distribution of X and Y is given by the pdf: $f(x, y) = \frac{1}{2\pi} e^{-(x^2+y^2)}$, $x \in (-\infty, \infty)$, $y \in (-\infty, \infty)$.

(a) Are X and Y independent? Why/why not? (2)

(b) Let $A = X + Y$, $B = X - Y$. Find $f(a, b)$ (6)

(c) Are A and B independent? Why/why not? (1)

(d) What is the distribution of A ? (1)

Pledge:

Question	Points	Score
1	15	
2	10	
3	10	
Total:	35	